

TOWARDS PLANE SPANNERS OF DEGREE 3*

Ahmad Biniaz,[†] Prosenjit Bose,[†] Jean-Lou De Carufel,[‡] Cyril Gavoille,[§] Anil Maheshwari,[†] and Michiel Smid[†]

ABSTRACT. Let S be a finite set of points in the plane. In this paper we consider the problem of computing plane spanners of degree at most three for S .

1. If S is in convex position, then we present an algorithm that constructs a plane $\frac{3+4\pi}{3}$ -spanner for S whose vertex degree is at most 3.
2. If S is the vertex set of a non-uniform rectangular lattice, then we present an algorithm that constructs a plane $3\sqrt{2}$ -spanner for S whose vertex degree is at most 3.
3. If S is in general position, then we show how to compute plane degree-3 spanners for S with a linear number of Steiner points.

1 Introduction

Let S be a finite set of points in the plane. A *geometric graph* is a graph $G = (S, E)$ with vertex set S and edge set E consisting of line segments connecting pairs of vertices. The *length* (or *weight*) of any edge (p, q) in E is defined to be the Euclidean distance $|pq|$ between p and q . The *length* of any path in G is defined to be the sum of the lengths of the edges on this path. For any two vertices p and q of S , their *shortest-path distance* in G , denoted by $|pq|_G$, is a minimum length of any path in G between p and q . For a real number $t \geq 1$, the graph G is a *t-spanner* of S if for any two points p and q in S , $|pq|_G \leq t|pq|$. The smallest value of t for which G is a *t-spanner* is called the *stretch factor* of G . A large number of algorithms have been proposed for constructing *t-spanners* for any given point set; see the book by Narasimhan and Smid [22].

The *degree* of a spanner is defined to be its maximum vertex degree. Note that 3 is a lower bound on the degree of a *t-spanner*, for every constant $t > 1$, because a Hamiltonian path through a set of points arranged in a grid has unbounded stretch factor (see [22] for more details). Even for points that are in convex position, 3 is a lower bound on the degree of a spanner (see Kanj *et al.* [18]).¹ Salowe [23] proved the existence of spanners of degree 4. Das and Heffernan [12] showed the existence of spanners of degree 3.

*A preliminary version of this paper appeared in the proceedings of the 27th International Symposium on Algorithms and Computation (ISAAC), 2016.

[†]Carleton University, ahmad.biniaz@gmail.com, {jit, anil, michiel}@scs.carleton.ca, supported by NSERC.

[‡]University of Ottawa, jdecaruf@uottawa.ca, supported by NSERC.

[§]University of Bordeaux, gavoille@labri.fr, supported by the ANR DESCARTES Project.

¹It can be shown that the stretch factor is $\Omega(\sqrt{n})$.

A *plane spanner* is a spanner whose edges do not cross each other. Chew [9] was the first to prove that plane spanners exist. Chew proved that the L_1 -Delaunay triangulation of a finite point set has stretch factor at most $\sqrt{10} \approx 3.16$ (observe that lengths in this graph are measured in the Euclidean metric). In the journal version [10], Chew proves that the Delaunay triangulation based on a convex distance function defined by an equilateral triangle is a 2-spanner. Dobkin *et al.* [13] proved that the L_2 -Delaunay triangulation is a t -spanner for $t = \frac{\pi(1+\sqrt{5})}{2} \approx 5.08$. Keil and Gutwin [19] improved the upper bound on the stretch factor to $t = \frac{4\pi}{3\sqrt{3}} \approx 2.42$. This was subsequently improved by Cui *et al.* [11] to $t = 2.33$ for the case when the point set is in convex position. Currently, the best result is due to Xia [24], who proved that t is less than 1.998. For points that are in convex position the current best upper bound on the stretch factor of plane spanners is 1.88 that was obtained by Amani *et al.* [1]. Regarding lower bounds, by considering the four vertices of a square, it is obvious that a plane t -spanner with $t < \sqrt{2}$ does not exist. Mulzer [21] has shown that every plane spanning graph of the vertices of a regular 21-gon has stretch factor at least 1.41611. Recently, Dumitrescu and Ghosh [15] improved the lower bound to 1.4308 for the vertices of a regular 23-gon.

The problem of constructing bounded-degree spanners that are plane and have small stretch factor has received considerable attention (e.g., see [4, 5, 6, 7, 8, 17, 18, 20]). Bonichon *et al.* [5] proved the existence of a degree 4 plane spanner with stretch factor 156.82. A simpler algorithm by Kanj *et al.* [18] constructs a degree 4 plane spanner with stretch factor 20; for points that are in convex position, this algorithm gives a plane spanner of degree at most 3 with the same stretch factor. Dumitrescu and Ghosh [14] considered plane spanners for uniform grids. For the infinite uniform square grid, they proved the existence of a plane spanner of degree 3 whose stretch factor is at most 2.607; the lower bound is $1 + \sqrt{2}$.

In this paper we consider bounded-degree plane spanners. In Section 3 we present an algorithm that computes a plane $\frac{3+4\pi}{3} \approx 5.189$ -spanner of degree 3 for points in convex position. In Section 4 we consider finite non-uniform rectangular grids; we present an algorithm that computes a degree 3 plane spanner whose stretch factor is at most $3\sqrt{2} \approx 4.25$. In Section 5 we show that any plane t -spanner for points in the plane that are in general position can be converted to a plane $(t + \epsilon)$ -spanner of degree at most 3 that uses a linear number of Steiner points, where $\epsilon > 0$ is an arbitrary small constant.

2 Preliminaries

For any two points p and q in the plane, let pq denote the line segment between p and q , let $\ell(p, q)$ denote the line passing through p and q , let $R(p \rightarrow q)$ denote the ray emanating from p and passing through q , and let $D(p, q)$ denote the closed disk that has pq as a diameter. Moreover, let $L(p, q)$ denote the lune of p and q , which is the intersection of the two closed disks of radius $|pq|$ that are centered at p and q .

Let S be a finite and non-empty set of points in the plane. We denote by $CH(S)$ the boundary of the convex hull of S . The *diameter* of S is the largest distance among the distances between all pairs of points of S . Any pair of points whose distance is equal

to the diameter is called a *diametral pair*. Any point of any diametral pair of S is called a *diametral point*. A *chain* is a sequence of points together with line segments connecting every pair of consecutive vertices.

Observation 1. Let S be a finite set of at least two points in the plane, and let $\{p, q\}$ be any diametral pair of S . Then, the points of S lie in $L(p, q)$.

The following theorem is a restatement of Theorem 7.11 in [3].

Theorem 1 (See [3]). If C_1 and C_2 are convex polygonal regions with $C_1 \subseteq C_2$, then the length of the boundary of C_1 is at most the length of the boundary of C_2 .

By simple calculations, one can verify the correctness of the following lemma; however, you can find a proof of it in [1].

Lemma 1. Let a, b , and c be three points in the plane, and let $\beta = \angle abc$. Then,

$$\frac{|ab| + |bc|}{|ac|} \leq \frac{1}{\sin(\beta/2)}.$$

Lemma 2. Let a and b be two points in the plane. Let c be a point that is on the boundary or in the interior of $L(a, b)$. Then, $\angle acb \geq \frac{\pi}{3}$.

Proof. Since $c \in L(a, b)$, we have $|ca| \leq |ab|$ and $|cb| \leq |ab|$. Thus, ab is a largest side of the triangle $\triangle abc$. This implies that $\angle acb$ is a largest internal angle of $\triangle abc$. Based on this, and since the sum of the internal angles of $\triangle abc$ is π , we conclude that $\angle acb \geq \frac{\pi}{3}$. \square

3 Plane Spanners for Points in Convex Position

In this section we consider degree-3 plane spanners for points that are in convex position. Let S be a finite set of points in the plane that are in convex position. Consider the two chains that are obtained from $CH(S)$ by removing any two edges. Let τ be the larger stretch factor of these two chains; notice that τ is not necessarily determined by the endpoints of the chain. In Section 3.1 we present an algorithm that computes a plane $(2\tau + 1)$ -spanner of degree 3 for S . Based on that, in Section 3.2 we show how to compute a plane $\frac{3+4\pi}{3}$ -spanner of degree 3 for S . Moreover, we show that if S is centrally symmetric, then there exists a plane $(\pi + 1)$ -spanner of degree 3 for S .

3.1 Spanner for Convex Double Chains

Let C_1 and C_2 be two chains of points in the plane that are separated by a straight line. Let S_1 and S_2 be the sets of vertices of C_1 and C_2 , respectively, and assume that $S_1 \cup S_2$ is in convex position. Let τ be a real number. In this section we show that if the stretch factor of each of C_1 and C_2 is at most τ , then there exists a plane $(2\tau + 1)$ -spanner for $S_1 \cup S_2$ whose degree is 3.




Figure 1: Illustration of the proof of Theorem 2.

In order to build such a spanner, we join C_1 and C_2 by a set of edges that form a matching. Thus, the spanner consists of C_1 , C_2 , and a set E of edges such that each edge has one endpoint in C_1 and one endpoint in C_2 . The set E is a matching, i.e., no two edges of E are incident on a same vertex. We show how to compute E recursively. Let (a, b) be the closest pair of vertices between C_1 and C_2 ; see Figure 1. Add this closest pair (a, b) to E . Then remove (a, b) from C_1 and C_2 , and recurse on the two pairs of chains obtained on each side of $\ell(a, b)$. Stop the recursion as soon as one of the chains is empty. Given C_1 and C_2 , the algorithm `MATCHING` computes a set E .

Algorithm 1 `MATCHING`(C_1, C_2)

Input: Two linearly separated chains C_1 and C_2 with the vertices of $C_1 \cup C_2$ in convex position.

Output: A matching between the points of C_1 and the points of C_2 .

- 1: **if** $C_1 = \emptyset$ or $C_2 = \emptyset$ **then**
 - 2: **return** \emptyset
 - 3: $(a, b) \leftarrow$ a closest pair of vertices between C_1 and C_2 such that $a \in C_1$ and $b \in C_2$
 - 4: $C'_1, C''_1 \leftarrow$ the two chains obtained by removing a from C_1
 - 5: $C'_2, C''_2 \leftarrow$ the two chains obtained by removing b from C_2
 - 6: **return** $\{ab\} \cup \text{MATCHING}(C'_1, C'_2) \cup \text{MATCHING}(C''_1, C''_2)$
-

In the rest of this section we prove the following theorem.

Theorem 2. Let $C_1 = (S_1, E_1)$ and $C_2 = (S_2, E_2)$ be two linearly separated chains of points in the plane, each with stretch factor at most τ , such that $S_1 \cup S_2$ is in convex position. Let E be the set of edges returned by algorithm `MATCHING`(C_1, C_2). Then, the graph $G = (S_1 \cup S_2, E_1 \cup E_2 \cup E)$ is a plane $(2\tau + 1)$ -spanner for $S_1 \cup S_2$ in which the degree of each of the endpoints of C_1 and C_2 is at most 2 and every other vertex has degree at most 3.

Proof. We prove this theorem by induction on $\min\{|S_1|, |S_2|\}$. As for the base cases, if $|S_1| = 0$, then $G = C_2$ is a plane τ -spanner whose vertex degree is at most 2. If $|S_2| = 0$, then $G = C_1$ is a plane τ -spanner whose vertex degree is at most 2.

Assume $|S_1| \geq 1$ and $|S_2| \geq 1$. Let ℓ be a line that separates C_1 and C_2 . Without loss of generality assume ℓ is horizontal, C_1 is above ℓ , and C_2 is below ℓ . Let (a, b) be the

pair of vertices selected by algorithm MATCHING, where (a, b) is a closest pair of vertices between C_1 and C_2 such that $a \in C_1$ and $b \in C_2$. Let C'_1 and C''_1 be the left and right sub-chains of C_1 , respectively, that are obtained by removing a ; see Figure 1. We obtain C'_2 and C''_2 similarly. Note that the chains C'_1 and C'_2 satisfy the conditions of Theorem 2. Let G' be the spanner obtained for the vertices of C'_1 and C'_2 . By the induction hypothesis, G' is a plane $(2\tau + 1)$ -spanner for the vertices of $C'_1 \cup C'_2$ in which the degree of each of the endpoints of C'_1 and C'_2 is at most 2 and every other vertex has degree at most 3. Similarly, let G'' be the spanner obtained for the vertices of C''_1 and C''_2 .

Observe that in G the degree of a , b , the right endpoint of C'_1 , the right endpoint of C'_2 , the left endpoint of C''_1 , and the left endpoint of C''_2 is at most 3. Moreover, in G , the degree of each endpoint of C_1 and C_2 is at most 2. Thus, G satisfies the degree condition. As for planarity, since G' and G'' are both plane and they are separated by $\ell(a, b)$, $G' \cup G''$ is also plane. Moreover, because of convexity, the edges that are incident on each of a and b do not cross any edge of $G' \cup G''$. Therefore, G is plane.

It remains to prove that the stretch factor of G is at most $2\tau + 1$. We are going to prove that for any two points $u, v \in S_1 \cup S_2$ we have $|uv|_G \leq (2\tau + 1)|uv|$. If both u and v belong to S_1 , or both belong to S_2 , then $|uv|_G \leq \tau|uv|$; this is valid because each of C_1 and C_2 has stretch factor at most τ . Assume $u \in S_1$ and $v \in S_2$. If $u, v \in G'$ or $u, v \in G''$ then, by the induction hypothesis, $|uv|_G \leq (2\tau + 1)|uv|$. Thus, it only remains to prove $|uv|_G \leq (2\tau + 1)|uv|$ for the following cases: (a) $u = a$ and $v \in C_2$, (b) $u \in C_1$ and $v = b$, (c) $u \in C'_1$ and $v \in C''_2$, and (d) $u \in C''_1$ and $v \in C'_2$. Because of symmetry we only prove cases (a) and (c).

First, we prove case (a). Assume $u = a$ and $v \in C_2$. Note that

$$|av|_G \leq |ab| + |bv|_{C_2} \leq |av| + \tau|bv|, \quad (1)$$

where the second inequality is valid since $|ab| \leq |av|$, by our choice of (a, b) , and since $|bv|_{C_2} \leq \tau|bv|$, given that the stretch factor of C_2 is at most τ . By the triangle inequality we have $|bv| \leq |ab| + |av|$. Since $|ab| \leq |av|$, we have $|bv| \leq 2|av|$. By combining this with Inequality (1) we get $|av|_G \leq (2\tau + 1)|av|$, which completes the proof for case (a).

Now, we prove case (c). Assume $u \in C'_1$ and $v \in C''_2$. Since S is in convex position, the polygon Q formed by u , a , v , and b is convex and its vertices appear in the order u, a, v, b . Note that

$$|uv|_G \leq |ua|_{C_1} + |ab| + |bv|_{C_2} \leq \tau|ua| + |uv| + \tau|bv| = |uv| + \tau(|ua| + |bv|), \quad (2)$$

where the second inequality is valid since $|ab| \leq |uv|$, by our choice of (a, b) , and since $|ua|_{C_1} \leq \tau|ua|$ and $|bv|_{C_2} \leq \tau|bv|$, given that the stretch factor of each of C_1 and C_2 is at most τ . Let c be the intersection point of ab and uv ; see Figure 1. By the triangle inequality, we have $|ua| \leq |uc| + |ca|$ and $|bv| \leq |bc| + |cv|$. It follows that $|ua| + |bv| \leq |uv| + |ab|$. Since $|ab| \leq |uv|$, we have $|ua| + |bv| \leq 2|uv|$. By combining this with Inequality (2) we get $|uv|_G \leq (2\tau + 1)|uv|$, which completes the proof of case (c). \square

3.2 Spanner for Points in Convex Position

In this section we show how to construct plane spanners of degree at most 3 for points that are in convex position.

Theorem 3. Let S be a finite set of points in the plane that is in convex position. Then, there exists a plane spanner for S whose stretch factor is at most $\frac{3+4\pi}{3}$ and whose vertex degree is at most 3.

The proof of this theorem uses the following result, which will be proved in Subsection 3.2.2:

Theorem 4. Let C be a convex chain with endpoints p and q . If C is in $L(p, q)$, then the stretch factor of C is at most $\frac{2\pi}{3}$.

Proof of Theorem 3. The proof is constructive; we present an algorithm that constructs such a spanner for S . The algorithm works as follows. Let (p, q) be a diametral pair of S . Consider the convex hull of S . Let C_1 and C_2 be the two chains obtained from $CH(S)$ by removing p and q (and their incident edges). Note that C_1 and C_2 are separated by $\ell(p, q)$. Let G' be the graph on $S \setminus \{p, q\}$ that contains the edges of C_1 , the edges of C_2 , and the edges obtained by running algorithm `MATCHING`(C_1, C_2). By Theorem 2, G' is plane and the endpoints of C_1 and C_2 have degree at most 2. We obtain a desired spanner, G , by connecting p and q , via their incident edges in $CH(S)$, to G' . This construction is summarized in algorithm `DEG3PLANESPANNER`.

Algorithm 2 `DEG3PLANESPANNER`(S)

Input: A non-empty finite set S of points in the plane that is in convex position

Output: A plane degree-3 spanner of S .

- 1: $(p, q) \leftarrow$ a diametral pair of S
 - 2: $C_1, C_2 \leftarrow$ the two chains obtained by removing p and q from $CH(S)$
 - 3: $E \leftarrow CH(S) \cup \text{MATCHING}(C_1, C_2)$
 - 4: **return** $G = (S, E)$
-

Observe that G is plane. Moreover, all vertices of G have degree at most 3; p and q have degree 2. Now we show that the stretch factor of G is at most $\frac{3+4\pi}{3} \approx 5.19$. Note that G consists of $CH(S)$ and a matching which is returned by algorithm `MATCHING`. Since p and q are diametral points, then by a result of [1], for any point $s \in S \setminus \{p\}$ we have $|ps|_{CH(S)} \leq 1.88|ps|$. Since $CH(S) \subseteq G$, we have $|ps|_G \leq 1.88|ps|$. By symmetry, the same result holds for q and any point $s \in S \setminus \{q\}$. Since (p, q) is a diametral pair of S , both C_1 and C_2 are in $L(p, q)$. Based on that, in Theorem 4, we will see that both C_1 and C_2 have stretch factor at most $\frac{2\pi}{3}$. Then, by Theorem 2, the stretch factor of G' is at most $\frac{3+4\pi}{3}$. Since $G' \subset G$, for any two points $r, s \in S \setminus \{p, q\}$ we have $|rs|_G \leq \frac{3+4\pi}{3}|rs|$. Therefore, the stretch factor of G is at most $\frac{3+4\pi}{3}$. This completes the proof of the theorem. \square

A point set S is said to be *centrally symmetric* (with respect to the origin), if for every point $p \in S$, the point $-p$ also belongs to S .

Theorem 5. Let S be a finite centrally symmetric point set in the plane that is in convex position. Then, there exists a plane spanner for S whose stretch factor is at most $\pi + 1$ and whose vertex degree is at most 3.

The proof of this theorem uses the following result, which will be proved in Subsection 3.2.1:

Theorem 6. Let C be a convex chain with endpoints p and q . If C is in $D(p, q)$, then the stretch factor of C is at most $\frac{\pi}{2}$.

Proof of Theorem 5. Let G be the graph obtained by $\text{DEG3PLANESPANNER}(S)$. As we have seen in the proof of Theorem 3, G is plane and has vertex degree at most 3. It remains to show that the stretch factor of G is at most $\pi + 1$. Let (p, q) be the diametral pair of S that is considered by algorithm DEG3PLANESPANNER . Since S is centrally symmetric, all points of S are in $D(p, q)$. Based on that, in Theorem 6, we will see that both C_1 and C_2 have stretch factor at most $\frac{\pi}{2}$. Then Theorem 2 implies that the stretch factor of G is at most $\pi + 1$. \square

In the following two subsections we will prove Theorems 4 and 6. Let C be a chain of points. For any two points u and v on C we denote by $\delta_C(u, v)$ the path between u and v on C . Recall that $|uv|_C$ denote the length of $\delta_C(u, v)$.

3.2.1 Proof of Theorem 6

Let p and q be the endpoints of the convex chain C . Assume that C is in $D(p, q)$. We are going to show that the stretch factor of C is at most $\frac{\pi}{2}$.

Since C is convex, it is contained in a half-disk of $D(p, q)$, i.e., a half-disk with diameter pq . Let u and v be any two points of C . We show that $\delta_C(u, v)$ is in $D(u, v)$. Then, by Theorem 1 the length of $\delta_C(u, v)$ is at most the length of the half-arc of $D(u, v)$, which is $\frac{\pi}{2}|uv|$. Without loss of generality assume that pq is horizontal, p is to the left of q , and C is above pq . Assume that u appears before v while traversing C from p to q . See Figure 2. We consider the following cases.




Figure 2: Proof of Theorem 6: the path $\delta_C(u, v)$ is inside the shaded regions, where (a) $u = p$ and $v \neq q$, (b) $u \neq p$, $v \neq q$, and $c \in D(p, q)$, and (c) $u \neq p$, $v \neq q$, and $c \notin D(p, q)$.

- $u = p$ and $v = q$. Then $\delta_C(p, q) = C$ is in $D(p, q)$ by the hypothesis.

- $u = p$ and $v \neq q$. Let v' be the intersection point of $R(q \rightarrow v)$ with the boundary of $D(p, q)$. See Figure 2(a). Observe that $\angle pv'v = \angle pv'q = \frac{\pi}{2}$. Thus, v' is on the boundary of $D(p, v)$. Since two circles can intersect in at most two points, p and v' are the only intersection points of the boundaries of $D(p, q)$ and $D(p, v)$. Thus, the clockwise arc $\widehat{pv'}$ on the boundary of $D(p, q)$ is inside $D(p, v)$. On the other hand, because of convexity, no point of $\delta_C(p, v)$ is to the right of $R(p \rightarrow v)$ or $R(q \rightarrow v)$. This implies that $\delta_C(p, v)$ is in $D(p, v)$.
- $u \neq p$ and $v = q$. The proof of this case is similar to the proof of the previous case.
- $u \neq p$ and $v \neq q$. Let c be the intersection point of $R(p \rightarrow u)$ and $R(q \rightarrow v)$. Because of convexity, $\delta_C(u, v)$ is in the triangle $\triangle ucv$. We look at two cases:
 - c is inside $D(p, q)$. In this case $\angle ucv \geq \frac{\pi}{2}$. See Figure 2(b). This implies that the point c , and consequently the triangle $\triangle ucv$, are inside $D(u, v)$. Thus, $\delta_C(u, v)$ is inside $D(u, v)$.
 - c is outside $D(p, q)$. Let u' (resp. v') be the intersection point of $R(p \rightarrow u)$ (resp. $R(q \rightarrow v)$) with $D(p, q)$. Because of the convexity and the fact that the path $\delta_C(u, v)$ is contained in $D(p, q)$, this path is inside the region that is bounded by line segments (u', u) , (u, v) , (v, v') and the boundary of $D(p, q)$; see the shaded region of Figure 2(c). Observe that by convexity $\angle uv'v > \angle pv'q = \frac{\pi}{2}$, and $\angle uu'v > \angle pu'q = \frac{\pi}{2}$. Thus, both u' and v' are inside $D(u, v)$. Consequently, the clockwise arc $\widehat{u'v'}$ on the boundary of $D(p, q)$ is inside $D(u, v)$. Therefore, $\delta_C(u, v)$ is inside $D(u, v)$.

3.2.2 Proof of Theorem 4

Let p and q be the endpoints of the convex chain C . Assume that C is in $L(p, q)$. We are going to show that the stretch factor of C is at most $\frac{2\pi}{3}$.

Since C is convex, it is contained in a half-lune of $L(p, q)$; i.e., a portion of $L(p, q)$ that is obtained by cutting it through pq . Let u and v be any two points of C . We show that the length of $\delta_C(u, v)$ is at most $\frac{2\pi}{3}$ times $|uv|$. Without loss of generality assume pq is horizontal, p is to the left of q , and C is above pq . Assume that u appears before v when traversing C from p to q . See Figure 3. We consider two cases: (1) $u = p$ or $v = q$, (2) $u \neq p$ and $v \neq q$.

1. $u = p$ or $v = q$. Without loss of generality assume $u = p$. If $v = q$ then $\delta_C(p, q)$ is inside the half-lune of $L(p, q)$ and by Theorem 1 the length of $\delta_C(p, q)$ is at most the length of the half-lune, which is $\frac{2\pi}{3}|pq|$. Assume $v \neq q$. For simplicity, we assume that $|pq| = 1$. Let $\alpha = \angle pqv$ and $x = |qv|$. If $\alpha = 0$, then the length of $\delta_C(p, v)$ is equal to $|pv|$ and we are done. Thus, assume that $\alpha > 0$. Moreover, since $v \neq q$, $x > 0$. We consider the following two cases: (a) $\alpha \leq \frac{\pi}{3}$, (b) $\alpha > \frac{\pi}{3}$.

- (a) $\alpha \leq \frac{\pi}{3}$. Let v' be the intersection point of $R(q \rightarrow v)$ and $L(p, q)$. Because of the convexity and the fact that the path $\delta_C(p, v)$ is contained in $L(p, q)$, this

path is inside the region that is bounded by line segments (p, v) , (v, v') and the boundary of $L(p, q)$; see the shaded region of Figure 3(a). By Theorem 1 we have the following inequality:

$$|pv|_C \leq |vv'| + |\widehat{pv'}| = 1 - x + \alpha,$$

where $|\widehat{pv'}|$ is the length of the clockwise arc from p to v' which has radius 1 and is centered at q . Note that

$$|pv| = \sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}.$$

Define

$$f(x, \alpha) = \frac{1 - x + \alpha}{\sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}}. \tag{3}$$

Then $\frac{|pv|_C}{|pv|} \leq f(x, \alpha)$. In Appendix A, we will show that $f(x, \alpha) \leq 2.04738 \leq \frac{2\pi}{3}$ for $0 < x \leq 1$ and $0 < \alpha \leq \frac{\pi}{3}$.




Figure 3: Proof of Theorem 4: the path $\delta_C(u, v)$ is inside the shaded regions, where (a) $u = p$ and $\alpha \leq \frac{\pi}{3}$, (b) $u = p$ and $\alpha > \frac{\pi}{3}$, (c) $u \neq p$, $v \neq q$, and $c \in L(p, q)$, and (d) $u \neq p$, $v \neq q$, and $c \notin L(p, q)$.

(b) $\alpha > \frac{\pi}{3}$. Let v' be the intersection point of $R(q \rightarrow v)$ and $L(p, q)$, and let r be the topmost intersection point of the two circles of radius $|pq|$ that are centered at

p and q , i.e., the highest point of $L(p, q)$. Because of the convexity and the fact that the path $\delta_C(p, v)$ is contained in $L(p, q)$, this path is inside the region that is bounded by line segments (p, v) , (v, v') and the boundary of $L(p, q)$; see the shaded region of Figure 3(b). By Theorem 1 we have the following inequality:

$$|pv|_C \leq |vv'| + |\widehat{rv'}| + |\widehat{pr}|,$$

where $|\widehat{rv'}|$ is the length of the clockwise arc from r to v' which has radius 1 and is centered at p and $|\widehat{pr}|$ is the length of the clockwise arc from p to r which has radius 1 and is centered at q . Let $\beta = \angle v'pq$. Then, $|\widehat{rv'}| = \frac{\pi}{3} - \beta$. Note that the triangle $\triangle pqv'$ is isosceles. Thus, $\beta = \pi - 2\alpha$ and $|qv'| = 2 \cos \alpha$. Thus $|vv'| = 2 \cos \alpha - x$. Therefore,

$$|pv|_C \leq (2 \cos \alpha - x) + \left(\frac{\pi}{3} - \beta\right) + \frac{\pi}{3} = 2(\alpha + \cos \alpha) - \left(x + \frac{\pi}{3}\right).$$

Define

$$g(x, \alpha) = \frac{2(\alpha + \cos \alpha) - (x + \pi/3)}{\sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}}. \tag{4}$$

Then $\frac{|pv|_C}{|pv|} \leq g(x, \alpha)$. In Appendix B, we will show that $g(x, \alpha) \leq \frac{2\pi}{3}$ for $0 < x \leq 2 \cos \alpha$ and $\frac{\pi}{3} \leq \alpha \leq \frac{\pi}{2}$.

2. $u \neq p$ and $v \neq q$. Let c be the intersection point of rays $R(p \rightarrow u)$ and $R(q \rightarrow v)$. We differentiate between two cases: (a) c is inside $L(p, q)$, (b) c is outside $L(p, q)$.

(a) c is inside $L(p, q)$. Because of the convexity, $\delta_C(u, v)$ is inside the triangle $\triangle uvc$; see Figure 3(c). Thus $|uv|_C \leq |uc| + |vc|$. Let $\beta = \angle ucv$. By Lemma 2 we have $\beta \geq \frac{\pi}{3}$. Based on this, and by Lemma 1, we have

$$\frac{|uv|_C}{|uv|} \leq \frac{|uc| + |vc|}{|uv|} \leq \frac{1}{\sin(\beta/2)} \leq \frac{1}{\sin(\pi/6)} = 2 < \frac{2\pi}{3}.$$

(b) c is outside $L(p, q)$. We reduce this case to case 1 where $u = p$ or $v = q$. See Figure 3(d). Let u' and v' be the intersection points of $R(p \rightarrow u)$ and $R(q \rightarrow v)$ with $L(p, q)$, respectively. Let δ_{uv} be the convex chain consisting of uu' , vv' , and the portion of the boundary of $L(p, q)$ that is between u' and v' (δ_{uv} is the bold chain in Figure 3(d)). Observe that the length of $\delta_C(u, v)$ is at most the length of δ_{uv} . Thus $\frac{|uv|_C}{|uv|} \leq \frac{|\delta_{uv}|}{|uv|}$. Let A be the convex region that is bounded by δ_{uv} and the segment (u, v) . Let A' be obtained from A by a homothetic transformation with respect to the center c and scale factor $\min\{\frac{|cp|}{|cu|}, \frac{|cq|}{|cv|}\}$. Assume $\frac{|cp|}{|cu|} \leq \frac{|cq|}{|cv|}$. Then, after this transformation u lies on p as shown in Figure 3(d). Let v'' be the point on qv where v ended up after this transformation. Let $\delta_{pv''}$ (the dashed chain in Figure 3(d)) be the chain obtained from δ_{uv} after this transformation. Since this transformation preserves ratios of distances we have $\frac{|\delta_{pv''}|}{|pv''|} = \frac{|\delta_{uv}|}{|uv|}$. Thus, $\frac{|uv|_C}{|uv|} \leq \frac{|\delta_{pv''}|}{|pv''|}$. Note that $\delta_{pv''}$ is in $L(p, q)$. To obtain an upper bound on $\frac{|\delta_{pv''}|}{|pv''|}$, we apply case 1 where $u = p$ and v'' plays the role of v .

4 Non-Uniform Rectangular Grid

In this section we build a plane spanner of degree three for the point set of the vertices of a non-uniform rectangular grid. In a finite non-uniform $m \times k$ grid, Λ , the vertices are arranged on the intersections of m horizontal and k vertical lines. The distances between the horizontal lines and the distances between the vertical lines are chosen arbitrary. The total number of vertices of Λ —the number of points of the underlying point set—is $n = m \cdot k$.




Figure 4: The grid Λ : (a) horizontal and vertical slabs, and (b) red and blue staircases.

If $m \in \{1, 2\}$ or $k \in \{1, 2\}$ then Λ is a plane spanner whose degree is at most 3 and whose stretch factor is at most $\sqrt{2}$. Assume $m \geq 3$ and $k \geq 3$. We present an algorithm that constructs a degree-3 plane spanner, G , for the points of Λ .

Let h_1, \dots, h_m be the horizontal lines of Λ from bottom to top. Similarly, let v_1, \dots, v_k be the vertical lines of Λ from left to right. For each pair (i, j) where $1 \leq i \leq m$ and $1 \leq j \leq k$, we denote by $p_{i,j}$ the vertex of Λ that is the intersection point of h_i and v_j . The vertices that are on h_1 , h_m , v_1 , or v_k are referred to as *boundary vertices*; other vertices are referred to as *internal vertices*. The edges both of whose endpoints are boundary vertices are referred to as *boundary edges*; other edges are referred to as *internal edges*. The grid Λ consists of $m - 1$ horizontal slabs (rows) and $k - 1$ vertical slabs (columns). Each horizontal slab H_i , with $1 \leq i < m$, is bounded by consecutive horizontal lines h_i and h_{i+1} . Each vertical slab V_j , with $1 \leq j < k$, is bounded by consecutive vertical lines v_j and v_{j+1} . See Figure 4(a). For each slab we define the *width* of that slab as the distance between the two parallel lines on its boundary.

We partition the set of internal edges of Λ into two sets: a set R of red edges and a

The graph G is plane because it is a subgraph of Λ . As for the degree constraint, since we add edges only between vertices that have degree 2 in $G' \cup E'$ (at most one edge per vertex), no vertex of degree 4 can appear. Thus, G has maximum degree 3. It only remains to show that G is a $3\sqrt{2}$ -spanner. Before that, we review some properties of G .

A *cell* of Λ refers to a region that is bounded by two consecutive horizontal lines and two consecutive vertical lines of Λ . Every face of G' is the union of the cells of Λ that are between two consecutive blue staircases and the boundary of Λ ; see Figure 5(a). Consider one iteration of the algorithm. We refer to the red edges that are not in $G' \cup E'$ as *missing edges*. For every face F in $G' \cup E'$ we define the set of *reflex vertices* of F as the vertices on the boundary of F that are incident on two missing edges in F ; reflex vertices have degree 2 in $G' \cup E'$. Notice that there exists a path of missing edges that connects the reflex vertices of F ; see Figure 5(a). Since the algorithm connects the reflex vertices by missing edges, all faces that have more than one reflex vertex have been broken into subfaces. Thus at the end of the algorithm every face of G contains at most one reflex vertex. Therefore, every face of G consists of one cell, two cells, or three cells. We refer to these faces as *1-cell*, *2-cell*, and *3-cell* faces, respectively. See the shaded faces in Figure 5(b).

Observation 2. Each face in G is either a 1-cell face, a 2-cell face, or a 3-cell face.

Lemma 3. Every missing edge that has an endpoint on the boundary of Λ is in a 2-cell face of G .

Proof. The proof is by contradiction. Assume (a, b) is a missing edge that has an endpoint on the boundary of Λ and lies in a 3-cell face F of G . Without loss of generality assume a is on the boundary. Thus, in G , b is the reflex vertex of F and has degree 2. At the moment the algorithm considers (a, b) , both a and b have degree 2, and thus, adds (a, b) to E' . Since all edges of E' , including (a, b) , are also edges of G , we get a contradiction with (a, b) being a missing edge. \square




Figure 6: Illustration of Lemmas 4 and 5: (a) (c, f) is the missing edge of a 2-cell face and f is a boundary vertex, (b) (c, f) is the missing edge of a 2-cell face and none of c and f are boundary vertices, and (c) (c, h) is a missing edge of a 3-cell face.

Lemma 4. Let (c, f) be the missing edge of a 2-cell face in G . Then, $|cf|_G \leq 3|cf|$.

Proof. Let $F = (a, b, c, d, e, f)$ be a 2-cell face of G with the edge (c, f) is missing. Without loss of generality assume that (c, f) is horizontal, f is to the left of c , and a, b, c, d, e, f is the

clockwise order of the vertices along the boundary of F ; see Figures 6(a) and 6(b). Note that $|cf| = |ab| = |de|$, $|af| = |bc|$, and $|ef| = |cd|$. We consider two cases:

- One of c and f is a boundary vertex. Without loss of generality assume that f is a boundary vertex, and thus, it has degree 2. See Figure 6(a). Then, (c, d) is a red edge that has been added during the algorithm. At the moment the algorithm considers (c, f) , the vertex c has degree 3, because otherwise, the algorithm adds (c, f) to E' , and hence to G . Thus, the edge (c, d) has been considered before (c, f) , which implies that $|cd| \leq |cf|$. Thus, the length of the path (c, d, e, f) , i.e., $|cd| + |de| + |ef|$, is at most $3|cf|$.
- Both c and f are internal vertices. See Figure 6(b). Then, (a, f) and (c, d) are red edges that have been added during the algorithm. At the moment the algorithm considers (c, f) , at least one of c and f has degree 3, because otherwise, the algorithm adds (c, f) to E' , and hence to G . Without loss of generality assume that c has degree 3. Thus, the edge (c, d) has been considered before (c, f) , which implies that $|cd| \leq |cf|$. Thus, the length of the path (c, d, e, f) , i.e., $|cd| + |de| + |ef|$, is at most $3|cf|$.

□

Lemma 5. Let (c, h) be a missing edge of a 3-cell face in G . Then, $|ch|_G \leq 3|ch|$.

Proof. Let $F = (a, b, c, d, e, f, g, h)$ be a 3-cell face in G with the edge (c, h) missing; see Figure 6(c). Let a, b, c, d, e, f, g, h be the clockwise order of the vertices along the boundary of F . Without loss of generality assume that c is the reflex vertex of F , the edge (c, h) is horizontal, and h is to the left of c . Note that $|ch| = |ab|$ and $|bc| = |ah|$. Since F is a 3-cell face, by Lemma 3, f is not a boundary vertex. Thus, (a, h) is a red edge that is added during the algorithm. At the moment the algorithm considers (c, h) , the vertex h has degree 3, because otherwise, the algorithm adds (c, h) to E' . Thus, (a, h) has been considered before (c, h) , which implies that $|ah| \leq |ch|$. Thus, the length of the path (c, b, a, h) , i.e., $|cb| + |ba| + |ah|$, is at most $3|ch|$. □




Figure 7: The path δ_{uv} is shown in dashed blue and the path δ'_{uv} is shown in bold blue.

Theorem 7. Let Λ be a finite non-uniform rectangular grid. Then, there exists a plane spanner for the point set of the vertices of Λ such that its degree is at most 3 and its stretch factor is at most $3\sqrt{2}$.


Proof. If Λ has less than three rows or less than three columns, then it is a plane $\sqrt{2}$ -spanner of degree 3. Assume Λ has at least three rows and at least three columns. Let G be the graph obtained by the algorithm described in this section. Then, G is plane and its vertex degree is at most 3. Since Λ is a $\sqrt{2}$ -spanner, between any two vertices u and v , there exists a path δ_{uv} in Λ such that $|\delta_{uv}| \leq \sqrt{2}|uv|$. By Lemmas 4 and 5, for any edge (a, b) on δ_{uv} that is not in G , there exists a path in G whose length is at most 3 times $|ab|$. Thus, δ_{uv} can be turned into a path δ'_{uv} in G such that $|\delta'_{uv}| \leq 3\sqrt{2}|uv|$; see Figure 7. Therefore, the stretch factor of G is at most $3\sqrt{2}$. \square

5 Final Remarks

In order to obtain plane spanners with small stretch factor, one may think of adding Steiner points² to the point set and build a spanner on the augmented point set. In the L_1 -metric, a plane 1-spanner of degree 4 can be computed by using $O(n \log n)$ Steiner points (see [16]). Arikati *et al.* [2] showed how to compute, in the L_1 -metric, a plane $(1 + \epsilon)$ -spanner with $O(n)$ Steiner points, for any $\epsilon > 0$. Moreover, for the Euclidean metric, they showed how to construct a plane $(\sqrt{2} + \epsilon)$ -spanner that uses $O(n)$ Steiner points and has degree 4.

Let S be a set of n points in the plane that is in general position; no three points are collinear. Let G be a plane t -spanner of S . We show how to construct, from G , a plane $(t + \epsilon)$ -spanner of degree 3 for S with $O(n)$ Steiner points, for any $\epsilon > 0$. Let CP denotes the closest pair distance in S and let $\epsilon' = \epsilon \cdot CP$.

For each point p of the point set S , consider a circle C_p with radius $\frac{\epsilon'}{\pi n}$ that is centered at p . Introduce a Steiner point on each intersection point of C_p with the edges of G that are incident on p . Also, introduce a Steiner point p' on C_p that is different from these intersection points. Delete the part of the edges of G inside each circle C_p (each edge $e = (p, q)$ of G turns into an edge e' of G' with endpoints on C_p and C_q). Add an edge from p to p' , and add a cycle whose edges connect consecutive Steiner points on the boundary of C_p . This results in a degree-3 geometric plane graph G' . For each vertex of degree k in G , we added $k + 1$ Steiner points in G' . Since G is planar, its total vertex degree is at most $6n - 12$. Thus, the number of Steiner points is $7n - 12$, in total (by removing Steiner vertices p' and b together with the edges incident on them in the above figure, keeping the edge (a, p) of G , and adding the edge (p, c) we get a different construction of G' that uses $5n - 12$ Steiner vertices).



A path δ_{uv} between two vertices u and v in G can be turned into a path δ'_{uv} in G' as follows. For each point p in S corresponding to an internal vertex of δ_{uv} incident on two edges e_1 and e_2 of δ_{uv} , replace the part of e_1 and e_2 inside C_p by the shorter of the two

²some points in the plane that do not belong to the input point set.

paths along C_p connecting the corresponding Steiner points. Also, for each point $p \in \{u, v\}$ incident on an edge e of δ_{uv} replace the part of e inside C_p with edge (p, p') together with the shorter of the two paths along C_p connecting p' and the Steiner point corresponding to e .

Since G is a t -spanner, $\frac{|\delta_{uv}|}{|uv|} \leq t$. Since the Steiner points are located at distance $\frac{\epsilon'}{\pi n}$ from points of S , the length of the detour caused by C_p at each vertex is at most $\frac{\epsilon'}{n}$. Since δ_{uv} has at most n vertices, the length of δ'_{uv} in G' is at most $|\delta_{uv}| + n \cdot \frac{\epsilon'}{n}$. Thus, $\frac{|\delta'_{uv}|}{|uv|} \leq \frac{|\delta_{uv}| + \epsilon'}{|uv|} = \frac{|\delta_{uv}|}{|uv|} + \epsilon \cdot \frac{CP}{|uv|} \leq t + \epsilon$, is valid because the closet pair distance CP is smaller than $|uv|$.

Acknowledgement

Some parts of this work have been done at the *Fourth Annual Workshop on Geometry and Graphs*, held at the Bellairs Research Institute in Barbados, March 6-11, 2016. The authors are grateful to the organizers and to the participants of this workshop. Also, we would like to thank Günter Rote for his comments that simplified the proofs of Theorems 4 and 6, and an anonymous referee for simplifying the construction of the spanner for non-uniform grids.

References

- [1] M. Amani, A. Biniiaz, P. Bose, A. Maheshwari, J. De Carufel, and M. Smid. A plane 1.88-spanner for points in convex position. *Journal of Computational Geometry*, 7(1):520–539, 2016.
- [2] S. R. Arikati, D. Z. Chen, L. P. Chew, G. Das, M. H. M. Smid, and C. D. Zaroliagis. Planar spanners and approximate shortest path queries among obstacles in the plane. In *Proceedings of the 4th European Symposium on Algorithms (ESA)*, pages 514–528, 1996.
- [3] R. V. Benson. *Euclidean geometry and convexity*. McGraw-Hill, 1966.
- [4] N. Bonichon, C. Gavoille, N. Hanusse, and L. Perkovic. Plane spanners of maximum degree six. In *Proceedings of the 37th International Colloquium on Automata, Languages and Programming (ICALP)*, pages 19–30, 2010.
- [5] N. Bonichon, I. A. Kanj, L. Perković, and G. Xia. There are plane spanners of degree 4 and moderate stretch factor. *Discrete & Computational Geometry*, 53(3):514–546, 2015.
- [6] P. Bose, P. Carmi, and L. Chaitman-Yerushalmi. On bounded degree plane strong geometric spanners. *Journal of Discrete Algorithms*, 15:16–31, 2012.
- [7] P. Bose, J. Gudmundsson, and M. H. M. Smid. Constructing plane spanners of bounded degree and low weight. *Algorithmica*, 42(3):249–264, 2005.

- [8] P. Bose, M. H. M. Smid, and D. Xu. Delaunay and diamond triangulations contain spanners of bounded degree. *International Journal of Computational Geometry and Applications*, 19(2):119–140, 2009.
- [9] L. P. Chew. There is a planar graph almost as good as the complete graph. In *Proceedings of the 2nd ACM Symposium on Computational Geometry*, pages 169–177, 1986.
- [10] L. P. Chew. There are planar graphs almost as good as the complete graph. *Journal of Computer and System Sciences*, 39:205–219, 1989.
- [11] S. Cui, I. A. Kanj, and G. Xia. On the stretch factor of Delaunay triangulations of points in convex position. *Computational Geometry: Theory and Applications*, 44:104–109, 2011.
- [12] G. Das and P. J. Heffernan. Constructing degree-3 spanners with other sparseness properties. *International Journal of Foundations of Computer Science*, 7(2):121–136, 1996.
- [13] D. P. Dobkin, S. J. Friedman, and K. J. Supowit. Delaunay graphs are almost as good as complete graphs. *Discrete & Computational Geometry*, 5:399–407, 1990.
- [14] A. Dumitrescu and A. Ghosh. Lattice spanners of low degree. *Discrete Mathematics, Algorithms and Applications*, 8(3):1–19, 2016.
- [15] A. Dumitrescu and A. Ghosh. Lower bounds on the dilation of plane spanners. *International Journal of Computational Geometry & Applications*, 26(2):89–110, 2016.
- [16] J. Gudmundsson, O. Klein, C. Knauer, and M. Smid. Small Manhattan networks and algorithmic applications for the earth mover’s distance. In *Proceedings of 23rd European Workshop on Computational Geometry (EuroCG)*, pages 174–177, 2007.
- [17] I. A. Kanj and L. Perkovic. On geometric spanners of Euclidean and unit disk graphs. In *Proceedings of the 25th Annual Symposium on Theoretical Aspects of Computer Science (STACS)*, pages 409–420, 2008.
- [18] I. A. Kanj, L. Perković, and D. Türkoğlu. Degree four plane spanners: Simpler and better. In *Proceedings of the 32nd International Symposium on Computational Geometry (SoCG)*, pages 45:1–45:15, 2016.
- [19] J. M. Keil and C. A. Gutwin. Classes of graphs which approximate the complete Euclidean graph. *Discrete & Computational Geometry*, 7:13–28, 1992.
- [20] X. Li and Y. Wang. Efficient construction of low weighted bounded degree planar spanner. *International Journal of Computational Geometry and Applications*, 14(1-2):69–84, 2004.
- [21] W. Mulzer. Minimum dilation triangulations for the regular n -gon. Master’s thesis, Freie Universität Berlin, Germany, 2004.

- [22] G. Narasimhan and M. Smid. *Geometric Spanner Networks*. Cambridge University Press, Cambridge, UK, 2007.
- [23] J. S. Salowe. Euclidean spanner graphs with degree four. *Discrete Applied Mathematics*, 54(1):55–66, 1994.
- [24] G. Xia. The stretch factor of the Delaunay triangulation is less than 1.998. *SIAM Journal on Computing*, 42:1620–1659, 2013.

A Upper bound for Function (3)

We prove that

$$f(x, \alpha) = \frac{1 - x + \alpha}{\sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}} \leq \sqrt{1 + \frac{1}{27} (3 + 2\pi)^2} \approx 2.04738$$

for all $0 < x \leq 1$, $0 < \alpha \leq \frac{\pi}{3}$. We rewrite $f(x, \alpha)$ as

$$\begin{aligned} \frac{1 - x + \alpha}{\sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}} &= \frac{1 - x\sqrt{\sin^2 \alpha + \cos^2 \alpha} + \alpha}{\sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}} \\ &= \frac{1 - \sqrt{(x \sin \alpha)^2 + (x \cos \alpha)^2} + \alpha}{\sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}} \\ &= \frac{1 - \sqrt{a^2 + b^2} + \alpha}{\sqrt{a^2 + (1 - b)^2}}, \end{aligned}$$

where $a = x \sin \alpha$ and $b = x \cos \alpha$. Note that $a^2 + (1 - b)^2 > 0$, because $\alpha > 0$. We prove the following equivalent inequality:

$$f(\alpha, a, b) = \frac{1 - \sqrt{a^2 + b^2} + \alpha}{\sqrt{a^2 + (1 - b)^2}} \leq \sqrt{1 + \frac{1}{27} (3 + 2\pi)^2} \approx 2.04738$$

for all $a = b \tan \alpha$, $0 < b \leq \cos \alpha$, $0 < \alpha \leq \frac{\pi}{3}$. We have

$$\frac{\partial}{\partial \alpha} f(\alpha, a, b) = \frac{1}{\sqrt{a^2 + (1 - b)^2}}.$$

Therefore, the system

$$\frac{\partial}{\partial \alpha} f(\alpha, a, b) = \frac{\partial}{\partial a} f(\alpha, a, b) = \frac{\partial}{\partial b} f(\alpha, a, b) = 0$$

does not have a solution. Thus, we look at the boundary conditions. Since $a = b \tan \alpha$, the boundary conditions are (1) $\alpha = 0$, (2) $\alpha = \frac{\pi}{3}$, (3) $b = 0$, and (4) $b = \cos \alpha$.

1. Since $a = b \tan \alpha$, we have

$$f(0, a, b) = f(0, 0, b) = 1 < \sqrt{1 + \frac{1}{27} (3 + 2\pi)^2}.$$

2. Since $a = b \tan \alpha$ and using elementary calculus, we can show that

$$\begin{aligned} f\left(\frac{\pi}{3}, a, b\right) &= f\left(\frac{\pi}{3}, \sqrt{3}b, b\right) = \frac{1 - 2b + \frac{\pi}{3}}{\sqrt{3b^2 + (1 - b)^2}} \\ &\leq f\left(\frac{\pi}{3}, \sqrt{3} \frac{\pi - 3}{4\pi + 6}, \frac{\pi - 3}{4\pi + 6}\right) = \sqrt{1 + \frac{1}{27} (3 + 2\pi)^2} \end{aligned}$$

for all $0 < b \leq \cos\left(\frac{\pi}{3}\right)$.

3. Since $a = b \tan \alpha$, we have

$$f(\alpha, a, 0) = f(\alpha, 0, 0) = 1 + \alpha \leq 1 + \frac{\pi}{3} < \sqrt{1 + \frac{1}{27} (3 + 2\pi)^2}$$

for all $0 < \alpha \leq \frac{\pi}{3}$.

4. Since $a = b \tan \alpha$ and using elementary calculus, we can show that

$$\begin{aligned} f(\alpha, a, \cos \alpha) &= f(\alpha, \sin \alpha, \cos \alpha) = \frac{\alpha}{\sqrt{2 - 2 \cos \alpha}} \\ &\leq f\left(\frac{\pi}{3}, \frac{\sqrt{3}}{2}, \frac{1}{2}\right) = \frac{\pi}{3} < \sqrt{1 + \frac{1}{27} (3 + 2\pi)^2} \end{aligned}$$

for all $0 < \alpha \leq \frac{\pi}{3}$.

B Upper bound for Function (4)

We prove that

$$g(x, \alpha) = \frac{2(\alpha + \cos \alpha) - (x + \frac{\pi}{3})}{\sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}} \leq \frac{2\pi}{3} \approx 2.09440$$

for all $0 < x \leq 2 \cos \alpha$, $\frac{\pi}{3} \leq \alpha \leq \frac{\pi}{2}$. We rewrite $g(x, \alpha)$ as

$$\begin{aligned} \frac{2(\alpha + \cos \alpha) - (x + \frac{\pi}{3})}{\sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}} &= \frac{2\left(\alpha + \frac{1}{\sqrt{\tan^2 \alpha + 1}}\right) - (x\sqrt{\sin^2 \alpha + \cos^2 \alpha} + \frac{\pi}{3})}{\sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}} \\ &= \frac{2\left(\alpha + \frac{1}{\sqrt{\left(\frac{x \sin \alpha}{x \cos \alpha}\right)^2 + 1}}\right) - \left(\sqrt{(x \sin \alpha)^2 + (x \cos \alpha)^2} + \frac{\pi}{3}\right)}{\sqrt{(x \sin \alpha)^2 + (1 - x \cos \alpha)^2}} \\ &= \frac{2\left(\alpha + \frac{1}{\sqrt{\left(\frac{a}{b}\right)^2 + 1}}\right) - (\sqrt{a^2 + b^2} + \frac{\pi}{3})}{\sqrt{a^2 + (1 - b)^2}} \\ &= \frac{2\left(\alpha + \frac{b}{\sqrt{a^2 + b^2}}\right) - (\sqrt{a^2 + b^2} + \frac{\pi}{3})}{\sqrt{a^2 + (1 - b)^2}} \\ &= \frac{6\left(\alpha + \frac{b}{\sqrt{a^2 + b^2}}\right) - (3\sqrt{a^2 + b^2} + \pi)}{3\sqrt{a^2 + (1 - b)^2}} \\ &= \frac{6\alpha - \pi}{3\sqrt{a^2 + (1 - b)^2}} - \frac{3\sqrt{a^2 + b^2} - \frac{6b}{\sqrt{a^2 + b^2}}}{3\sqrt{a^2 + (1 - b)^2}} \\ &= \frac{6\alpha - \pi}{3\sqrt{a^2 + (1 - b)^2}} - \frac{a^2 + (1 - b)^2 - 1}{\sqrt{a^2 + b^2} \sqrt{a^2 + (1 - b)^2}}, \end{aligned}$$

where $a = x \sin \alpha$ and $b = x \cos \alpha$. Note that $a^2 + b^2 > 0$ and $a^2 + (1 - b)^2 > 0$, because $x > 0$ and $\alpha > 0$. We prove the following equivalent inequality:

$$g(\alpha, a, b) = \frac{6\alpha - \pi}{3\sqrt{a^2 + (1 - b)^2}} - \frac{a^2 + (1 - b)^2 - 1}{\sqrt{a^2 + b^2}\sqrt{a^2 + (1 - b)^2}} \leq \frac{2\pi}{3} \approx 2.09440$$

for all $a = b \tan \alpha$, $0 \leq b \leq 2 \cos^2 \alpha$, $\frac{\pi}{3} \leq \alpha \leq \frac{\pi}{2}$. We have

$$\frac{\partial}{\partial \alpha} g(\alpha, a, b) = \frac{2}{\sqrt{a^2 + (1 - b)^2}}.$$

Therefore, the system

$$\frac{\partial}{\partial \alpha} g(\alpha, a, b) = \frac{\partial}{\partial a} g(\alpha, a, b) = \frac{\partial}{\partial b} g(\alpha, a, b) = 0$$

does not have a solution. Thus, we look at the boundary conditions. Since $a = b \tan \alpha$, the boundary conditions are (1) $\alpha = \frac{\pi}{3}$, (2) $\alpha = \frac{\pi}{2}$, (3) $b = 0$, and (4) $b = 2 \cos^2 \alpha$.

1. Since $a = b \tan \alpha$ and using elementary calculus, we can show that

$$\begin{aligned} g\left(\frac{\pi}{3}, a, b\right) &= g\left(\frac{\pi}{3}, \sqrt{3}b, b\right) = \frac{1 - 2b + \frac{\pi}{3}}{\sqrt{3b^2 + (1 - b)^2}} \\ &\leq g\left(\frac{\pi}{3}, \sqrt{3} \frac{\pi - 3}{4\pi + 6}, \frac{\pi - 3}{4\pi + 6}\right) = \sqrt{1 + \frac{1}{27} (3 + 2\pi)^2} < \frac{2\pi}{3} \end{aligned}$$

for all $0 \leq b \leq 2 \cos^2(\frac{\pi}{3})$.

2. When $\alpha = \frac{\pi}{2}$, $a = b \tan \alpha$ is not well-defined. Instead, we write $b = a \cot \alpha = 0$ and $0 \leq a \leq 2 \sin \alpha \cos \alpha = 0$, from which $a = b = 0$. We have

$$g\left(\frac{\pi}{2}, 0, 0\right) = \frac{2\pi}{3}.$$

3. Since $a = b \tan \alpha$, we have

$$g(\alpha, a, 0) = g(\alpha, 0, 0) = \frac{6\alpha - \pi}{3} \leq \frac{2\pi}{3}$$

for all $\frac{\pi}{3} \leq \alpha \leq \frac{\pi}{2}$.

4. Since $a = b \tan \alpha$, we have

$$g\left(\alpha, a, 2 \cos^2 \alpha\right) = g\left(\alpha, 2 \sin \alpha \cos \alpha, 2 \cos^2 \alpha\right) = 2\alpha - \frac{\pi}{3} \leq \frac{2\pi}{3}$$

for all $\frac{\pi}{3} \leq \alpha \leq \frac{\pi}{2}$.