

On The Colouring Problem In The Physical Local Model

Cyril GAVOILLE **Ghazal KACHIGAR** Gilles ZÉMOR

Institut de Mathématiques de Bordeaux

LaBRI

October 2, 2017


Introduction

Distributed protocol

Centralised protocol



Distributed protocol



A distributed protocol may use :

- **no randomness:** $\mathbb{P}(y_i^*|x_i^*) = 1$, $\mathbb{P}(y_i^*|x_i) = 0$ for all $x_i \neq x_i^*$.
- **local randomness:** $\mathbb{P}(y_1, \dots, y_n|x_1, \dots, x_n) = \prod_{i=1}^n \mathbb{P}(y_i|x_i, \lambda_i)$.
- **shared randomness:** $\mathbb{P}(y_1, \dots, y_n|x_1, \dots, x_n) = \sum_{\lambda} \mathbb{P}(\lambda) \prod_{i=1}^n \mathbb{P}(y_i|x_i, \lambda)$.
- **quantum entanglement**


Introduction

CHSH game

[Bell 64] : Existence of correlation arising from quantum mechanics that cannot be modelled by a "local hidden variable theory", i.e.

"shared randomness < quantum entanglement"

CHSH game



Winning condition :

$$y_a \oplus y_b = x_a \wedge x_b$$

Probability of winning :

- Using shared randomness : at most 0.75.
- Using a quantum "Bell state" : $\cos^2(\pi/8) \approx 0.86$.

Introduction

Non-Signaling condition

Correlations arising from the quantum solution are **non-signaling**, i.e. the output of Alice doesn't give any information on the input of Bob and vice-versa.

Mathematically :

$$\sum_{y_b} \mathbb{P}(y_a, y_b | x_a, x_b) = \sum_{y_b} \mathbb{P}(y_a, y_b | x_a, x'_b) = \mathbb{P}(y_a | x_a)$$

and

$$\sum_{y_a} \mathbb{P}(y_a, y_b | x_a, x_b) = \sum_{y_a} \mathbb{P}(y_a, y_b | x'_a, x_b) = \mathbb{P}(y_b | x_b)$$

We have

Classical \subsetneq Quantum \subsetneq Non-Signaling

\Rightarrow Not Non-Signaling implies not Quantum

\Rightarrow [Arfaoui 14] showed that for 2 players with binary input and output and output condition $\neq y_a \oplus y_b$ the best non-signaling probability distribution is classical.

Introduction

LOCAL model


Suppose we have a graph $G = (V, E)$ modelling a communication network.

LOCAL model

- Every node has a (unique) **identifier**.
- One **round** of communication: send & receive information to neighbours & do computation.
- k rounds of communication \Leftrightarrow exchange with neighbours at distance $\leq k$ and do computation
- "Infinite" local computing power and bandwidth.

Introduction

Colouring Problem



Distributed Colouring Problem in the LOCAL model

How many rounds of communication are necessary and sufficient for q -colouring a graph ?


E.g. $q = \Delta + 1$ and graph=cycle or path.

[**Cole & Vishkin 86**] : $\log^*(n)$ rounds of communication are sufficient.

[**Linial 92**] : $\log^*(n)$ rounds of communication are necessary.


Physical Locality

[Gavoille, Kosowki & Markiewicz 09] : Non-Signaling + LOCAL = ϕ -local
Non-Signaling




ϕ -local

Input



Output



Physical Locality

An important property

[Barrett, Noah Linden, Massar, Pironio, Sandu, Popescu & Roberts 05] : if the non-signaling property is satisfied for a coalition of $n - 1$ players, then it is satisfied for any sub-coalition of $k < n - 1$ players.

$$\begin{aligned}\sum_{y_2, y_3} \mathbb{P}(y_1, y_2, y_3 | x_1, x_2, x_3) &= \sum_{y_2} \sum_{y_3} \mathbb{P}(y_1, y_2, y_3 | x_1, x_2, x'_3) \\ &= \sum_{y_3} \sum_{y_2} \mathbb{P}(y_1, y_2, y_3 | x_1, x'_2, x'_3) \\ &= \sum_{y_2, y_3} \mathbb{P}(y_1, y_2, y_3 | x_1, x'_2, x'_3)\end{aligned}$$

If a coalition of $m \leq n - 1$ players such that there is more than one global output corresponding to their inputs satisfies ϕ -local, so does any sub-coalition of $k < m$ players.

⇒ **To check for non-signaling/ ϕ -local**, need only look at biggest coalitions.

⇒ **To show non-signaling/ ϕ -local isn't satisfied**, small coalitions are sufficient.

Physical Locality

An example

Example : 2-colouring in an undirected n -path is ϕ -local($\lfloor \frac{n}{3} \rfloor$).


E.g. $n = 9$, $\lfloor \frac{9}{3} \rfloor = 3$




Physical Locality

A formal definition


We define the k -neighbourhood $\mathcal{N}_k(v)$ of a vertex v as the set of vertices at distance less than or equal to k .



$\mathcal{N}_0(2)$



$\mathcal{N}_1(2)$



$\mathcal{N}_2(2)$

Definition

Let $G = (E, V)$ be a directed or undirected graph and let $(X_v)_{v \in V}$ be a stochastic process. $(X_v)_{v \in V}$ is said to be ϕ -local(k) if, for every $m \leq |V|$ and sets of vertices $S = \{s_1, \dots, s_m\}$ and $T = \{t_1, \dots, t_m\}$ such that the graphs induced by $S \cup \mathcal{N}_k(S)$ and $T \cup \mathcal{N}_k(T)$ are isomorphic, we have

$$\mathbb{P}(X_S) = \mathbb{P}(X_T)$$

Physical Locality

Links to probability theory

Let $(X_n)_{n \in \mathbb{Z}}$ be a stochastic process on \mathbb{Z} .

(1) "Radius" of the information necessary to compute the value of a X_n ?

r-block factor

$(X_n)_{n \in \mathbb{Z}}$ is *r*-block factor of an iid process $(Y_n)_{n \in \mathbb{Z}}$ if $X_n = f(Y_n, \dots, Y_{n+r-1})$ for every n .

\Rightarrow Distributed computability in LOCAL model.

(2) "Radius" beyond which no information on the value of X_n escapes?

k-dependence

$(X_n)_{n \in \mathbb{Z}}$ is *k*-dependent if the distributions $(X_{\leq n})$ and $(X_{\geq m})$ are independent for every n, m with $|m - n| > k$.

Question : is *k*-dependence the same as ϕ -local(ℓ) for some ℓ ?

Physical Locality

Some results on k -dependence

It is easy to verify that: k -dependent \Rightarrow directed ϕ -local(k), undirected ϕ -local($\lfloor k/2 \rfloor$).
[Holroyd & Liggett 15], [Holroyd & Liggett 16] proved the following

k -dependent colouring of \mathbb{Z}

There exists a 1-dependent q -colouring process for every $q \geq 4$ and a 2-dependent 3-colouring process but no 1-dependent 3-colouring process.

Directed ϕ -local(1) 4-colouring and undirected ϕ -local(1) or directed ϕ -local(2) 3-colouring of the infinite path is possible.

1-dependent colouring of other graphs

For the following graphs G , the least possible number q of colours such that G admits a 1-dependent colouring is:

- $G = \mathbb{Z}^2$: 9
- $G = \mathbb{Z}^3$: 12
- $G =$ infinite Δ -regular tree, $\Delta \geq 2$: $\frac{(\Delta-1)^{\Delta-1}}{\Delta^\Delta}$

Physical Locality

k -dependence and ϕ -locality

Question

is k -dependence the same as ϕ -local(ℓ) for some ℓ ?

We studied this question by looking at the colouring problem on the path graph.

Our results:

- In general, no.
- In the case of the colouring problem: not exactly, but beyond a certain value of n , ϕ -local(1) 3-colouring of a directed n -path is not possible.

Our results

Idea : given a graph $G = (E, V)$ along with a q -colouring, choose a colour q^* and replace all its occurrences by 1 and put 0 everywhere else. This induces a binary process $(J_v)_{v \in V}$ which is k -dependent if the original colouring is.



$p^* = \sup\{p, \exists \text{ 1-dependent binary process s.t. for every } v \in V \mathbb{P}(J_v = 1) = p\}$, then $q \geq \frac{1}{p^*}$.

Theorem [Holroyd & Liggett 2016]

Let $G = (V, E)$ be a graph, p^* as above and $p \in [0, 1]$. We have $p \leq p^*$ iff $Z_A(-p) \geq 0$ for every finite $A \subset V$, where Z_A is the independence polynomial of the induced subgraph of A .

Thus on the n -path, $p^* < 1/3$ as soon as $n = 5$ and $\lim_{n \rightarrow \infty} p^* = \frac{1}{4}$.

Our results

Define $p_1 = \mathbb{P}(1)$, $p_2 = \mathbb{P}(1 * 1)$, $p_i = \mathbb{P}((1*)^i)$.

Catalan numbers : $c_n = \frac{1}{n+1} \binom{2n}{n}$, in particular $\frac{c_n}{c_{n+1}} = \frac{n+2}{2(2n+1)}$

For the directed path graph of length $n = 2\ell$ or $n = 2\ell + 1$ in the ϕ -local(1) model

$$- p_1 \leq \frac{c_\ell}{c_{\ell+1}}$$

$$- p_2 \leq \frac{c_{\ell-1}}{c_\ell} p_1 \leq \frac{c_{\ell-1}}{c_{\ell+1}}$$

$$- p_i \leq \frac{c_{\ell-i+1}}{c_{\ell-i}} p_{i-1} \leq \frac{c_{\ell-i+1}}{c_{\ell+1}}$$

Thus $p_1 < \frac{1}{3}$ as soon as $\ell = 5$, i.e. $n = 10$.

Let $p_i^* = \sup(p_i)$, then $\lim_{n \rightarrow \infty} p_1^* = \frac{1}{4}$ and $\lim_{n \rightarrow \infty} p_i^* = \frac{1}{4^i} = (p_1^*)^i$.

Our results

Method

The probabilities of each pattern is a linear function of the p_i .

Goal : maximise p_1 such that every probability is between 0 and 1.

Maximise $c^T p$ under the constraints $A p \leq b$

Minimise $b^T q$ under the constraints $A^T q \leq c$

Duality theorem : objective function has same value for optimal solutions of dual and primal problems.

We get $p_1 \leq \frac{c_\ell}{c_{\ell+1}} \Rightarrow$ remove first line of A , rearrange and solve again for p_2 , etc.

Summary

- ϕ -local model useful for finding if there might be a quantum-classical difference in number of rounds in distributed graph algorithms.
- k -dependence from probability theory is consistent with ϕ -local but stronger.
- $\log^*(n)$ classical rounds necessary and sufficient for solving the distributed graph colouring problem.
- There is a 1-dependent q -colouring probability distribution for $q \geq 4$ and a 2-dependent 3-colouring probability distribution on the path graph.
- Can barely do better in ϕ -local.
- Not easy to study for other families of graphs.

Thank You



For Your Attention